



Post-Operative Instructions Great Toe Chielectomy

Paul Cammack, MD
763-302-2226

WOUND CARE

- You will be placed into a soft dressing at the time of surgery. This dressing should remain in place and be kept clean and dry for the first 48 hours after surgery.
- You may remove your dressing 48 hours after surgery, and then you may get your incision wet in the shower.
- Re-dress your incision with dry gauze as needed if it is draining.
- You will be given a post-op shoe to use after surgery. You should use this shoe for walking until your sutures are removed.
- At your first post-op visit, your sutures will be removed, and Steri-Strips will be placed if your skin is healed.
- Steri-Strips will fall off in a few days. If they have not fallen off 5 days after your sutures are removed, you should remove them to allow air to the incision.
- Do not soak your incision in a tub, pool, hot tub, or lake until the incision is fully healed.
- Do not put lotions, creams, or ointments directly on the incision until it is fully healed.

ACTIVITY

- You should elevate your foot as much as possible after surgery. This helps reduce swelling and promote healing.
- You should ice your foot for 20 minutes at least twice a day until your follow-up appointment.
- You may weight bear as tolerated on your foot immediately after surgery.
- You may use crutches or a walker as needed for comfort, although they are not required.
- You may drive when you are comfortable and are no longer taking narcotic pain medications.
- Return to work will be discussed at your post-op appointment and will depend on your job requirements.

DIET

- Anesthetic medications and pain medicines can cause nausea and constipation. It is wise to begin with liquids and bland foods until any nausea has resolved. If you have persistent nausea and vomiting, call the office.

- Be sure to drink plenty of fluids as you are recovering, especially while taking pain medicine. This will reduce the risk of constipation.
- An over-the-counter stool softener is also advised while taking narcotic pain medicine.
- Smoking severely inhibits healing, particularly after ankle or foot surgery. **DO NOT SMOKE** while you are recovering from your foot procedure.

MEDICATION

- You may resume your regular pre-operative medications the evening of surgery unless otherwise directed.
- You will be prescribed narcotic pain medicine after surgery. Use as directed. Usually this will be needed for several days to weeks after surgery. This medicine has side effects and can be addicting. Try and wean off the narcotics as soon as you can. In most cases, patients should be off all narcotic pain medicine 6 weeks after surgery.
- You likely had a nerve block at the time of your surgery. This helps with post-op pain but will wear off 12 – 24 hours after your surgery. Don't wait until the block is fully worn off to start taking medicine. It is wise to have some medicine on board before the block wears off.
- Over-the-counter medicines such as Tylenol and ibuprofen are helpful. You should alternate these at the prescribed doses along with your prescribed medication.
- Pain medication will help with but not eliminate pain after surgery. Some discomfort is to be expected.

BLOOD CLOTS

- Blood clots, also known as deep vein thrombosis or DVT, are an uncommon occurrence after ankle or foot surgery but can be dangerous, even life threatening.
- Wiggling your toes and moving around help prevent blood clots.
- You will also be instructed to take medication, likely daily aspirin, to help prevent blood clots. You should stay on your aspirin until you are walking without your post-op shoe.
- Tell your surgeon if you have a personal or family history of blood clots.
- Women on oral contraceptives have an increased risk of blood clots.
- If your calf or leg becomes increasingly swollen and tender, call the office.
- If you develop pain in your chest or shortness of breath, call 911.

FOLLOW-UP

- Your first 2 follow-up appointments are scheduled when your surgery is scheduled. Your first appointment should be roughly 2 weeks after surgery. If you don't have an appointment, call the office.
- Call the office if you have any of the following:
 - Fever greater than 101.5 degrees
 - Pain that is not controlled with medication
 - Persistent nausea or vomiting
 - Worsening numbness or tingling in your toes (after your block wears off)

- Worsening redness or drainage from your incision
- Any other concerns