



Jozef Murar, M.D.

TCO Edina – Crosstown
 4010 W 65th St, Edina, MN 55435
 Tel: 952-456-7000 Fax: 952-832-0477
www.tcomn.com

Post-Operative Rehabilitation Prescription for:
Arthroscopic Anterior/Posterior Shoulder Stabilization: (M24.819)

Patient Name: _____ **Date:** _____
Treatment Frequency: 2 – 3 times per week **Duration of Therapy Prescription:** 8 weeks

Phase I MAXIMAL PROTECTION (0 - 4 Weeks Post-Op)

- Immobilization** for 4 weeks using sling.
- Elbow A/AAROM: flexion and extension.
- Protect anterior/posterior capsule from stretch**
- Limit FE (forward elevation in the scapular plane) to 110 degrees
- Limit ER (external rotation) to neutral 0 degrees
- Pendulums permitted in sling.
- Modalities (i.e. CryoCuff) PRN(as needed).
- Wrist and gripping exercises.
- Sub-maximal / Deltoid **isometrics** (Except internal rotation secondary to subscapularis reattachment.)
- Removal of sling for showering: **maintain arm in sling position.**

Phase II MODERATE PROTECTION (4 - 6 Weeks Post-Op)

- At 4 weeks PROM: pulley for flexion, pendulum exercises. **Progress as tolerated.**
- A/AAROM Limit FE (forward elevation in the scapular plane) to 140 degrees
- A/AAROM Limit ER (external rotation) to 45 degrees
- Progress from AAROM to AROM:
 - 1) Quality movement only-**avoid forcing** active motion with substitution patterns.
 - 2) Remember the effects of gravity on the limb, do gravity eliminated motions first i.e. Supine flexion.
- Deltoid isometrics.
- Lightly resisted **elbow** flexion.
- Continue with wrist exercises
- Modalities PRN.
- Discontinue sling at 4-6 weeks.

Phase III MINIMAL PROTECTION/MILD STRENGTHENING (6 - 12 Weeks)

- ❑ A/AAROM No Limit FE (forward elevation in the scapular plane)
- ❑ A/AAROM No Limit ER (external rotation)
- ❑ 10-12 weeks, AIAA/PROM to improve ER with arm in **45 degree** abduction.
- ❑ AROM all directions below horizontal, light resisted motions in all planes.
- ❑ AROM activities to restore flexion, IR, horiz ADD.
- ❑ Deltoid, Rotator Cuff isometrics progressing to isotonic.
- ❑ PRE's for scapular muscles, latissimus, biceps, triceps.
- ❑ PRE's work rotators in isolation (use modified neutral).
- ❑ Joint mobilization (posterior glides).
- ❑ Emphasize **posterior cuff, latissimus, and scapular muscle** strengthening, stressing eccentrics.
- ❑ Utilize **exercise arcs** that protect anterior capsule from stress during PRE's.
- ❑ Keep **all** strength exercises below the **horizontal plane** in this phase.

Phase IV STRENGTHENING (12 - 16 Weeks Post-Op)

- ❑ CRITERIA:
 - 1) Pain-free AROM
 - 2) Pain-free with manual muscle test
 - 3) Progress by response to treatment
- ❑ AROM activities to restore full ROM.
- ❑ Restore scapulohumeral rhythm.
- ❑ Joint mobilization.
- ❑ Aggressive scapular stabilization and eccentric strengthening program.
- ❑ Initiate isotonic shoulder strengthening exercises including: side lying ER, prone arm raises at 0, 90, 120 degrees, elevation in the plane of the scapula with IR and ER, lat pulldown close grip, and prone ER.
- ❑ Dynamic stabilization WB and NWB.
- ❑ PRE's for all upper quarter musculature (begin to integrate upper extremity patterns). Continue to emphasize eccentrics and glenohumeral stabilization.
- ❑ All PRE's are below the horizontal plane for non-throwers.
 - 1) Begin isokinetics.
 - 2) Begin muscle endurance activities (UBE).
 - High seat and low resistance
 - Must be able to do active shoulder flexion to 90 degrees without substitution
 - 3) Continue with agility exercises.
 - 4) Advanced functional exercises.
 - 5) Isokinetic test.
 - 6) Functional test assessment.
 - 7) Full return to sporting activities.

Please send progress notes.

Physician's Signature: _____

(I have medically prescribed the above treatments)

Jozef Murar, M.D.

Orthopaedic Surgeon