

SHOULDER CAPSULAR SHIFT / BANKART-LABRAL REPAIR

Procedure Name: _____

Date: _____

Anesthesia: General Sedation Block _____

- You have received sedation; rest and relax the day of surgery. Please be aware of possible dizziness and exercise caution when you are up. **A responsible adult must be with the patient for 24 hours following surgery for safety and falls prevention.**
- Begin with liquids, and advance diet as tolerated; avoid greasy and spicy foods.
- No important decisions or signing of legal documents for 24 hours. Do not operate power machinery.
- Do not drive for 24 hours following surgery or while taking narcotic pain medications.
- No alcoholic beverages for 24 hours, or while taking narcotics.
- If you are unable to urinate, feel uncomfortable, and it has been 8-10 hours since you last urinated, go to an urgent care or the ER.
- Nerve Block** If you have had a nerve block, be sure to start taking your pain medication as you feel sensation returning to your operative extremity. Typically, a nerve block will last between 8-24 hours from the time of injection. Notify your surgeon if the nerve block lasts more than 48 hours. Do not use your operative extremity until the nerve block has worn off.

*****In the event of a medical emergency, dial 911 or go to the nearest emergency department**

Report to your Surgeon or visit our Twin Cities Orthopedics Urgent Care in Waconia, Excelsior, or Edina (Sun-Sat 8am-8pm) for any of the following:

- Fever over 101 degrees for more than 24 hours
- Foul drainage, redness, or warmth at the operative site
- Large amount of bleeding or drainage
- Severe and uncontrolled pain
- Persistent nausea or vomiting
- Hives, rash, or intolerance (Stop medications)
- Redness, warmth, or extreme pain in the calf

POST-OP SHOULDER SUPPORT:

Always use your sling, day and night for the first two weeks. You may remove the sling to shower or for exercises. During that time, you may bring the arm completely down to the side temporarily.

SLEEPING:

After your shoulder surgery, you may sleep more easily in a recliner or propped up on several pillows in a semi-sitting position. This is commonly noted for four weeks following surgery. You may sleep flat on your back when you feel comfortable.

MOTION RESTRICTIONS:

You can reach to shoulder height only after two weeks. The arm should not be rotated beyond the side of the body for the first six weeks. This is 0 degrees of external rotation. You may flex the shoulder to 90 degrees by six weeks.

DRESSING:

- Change dressing daily beginning second day post-op. Please leave steri strips in place if possible. Re-dress with a light gauze covering. You may leave the wound uncovered when dry if you wish. You may shower 48 hours after surgery.

- Be alert for signs of possible infection: redness, heat, red streaks, or elevated temperature (over 101 degrees F). Contact your doctor if any of these occur.

PAIN CONTROL:



- Apply ice to the surgical area to reduce swelling and pain for 24-72 hours following the procedure. Do not place ice directly on skin or it may cause a burn. Wrap a towel around an ice pack and apply to the surgical site for no more than 20 minutes at a time per hour.
- Call your pharmacy for prescription refills. Prescription refills are not ordered over the weekend or after hours.
- **DO NOT DRIVE WHILE TAKING NARCOTIC PAIN MEDICATIONS** and until instructed by your physician.

POST-OPERATIVE DOCTOR'S APPOINTMENT:

Please call (952) 442-2163 to make your appointment for 10-14 days post-op.

EXERCISES:

You may begin shoulder mobility exercises day two post-op. See below for descriptions and illustrations of your exercises.

<p>Active range of motion for elbow:</p>	<p>Begin with arm straight. Grasp arm at wrist. Move elbow through available range and return to start position. Special Instructions: Use a firm, yet soft grip, cradling arm. Do not squeeze over bony areas. Perform 3 sets of 30 repetitions, three times a day. Rest 1 minute between sets. Perform 1 repetition every 4 seconds.</p>	
<p>Shoulder blade squeezes</p>	<p>Stand with arms at sides. Squeeze both shoulder blades together. Relax and repeat. Perform 3 sets of 20 repetitions, three times a day. Rest 1 minute between sets. Perform every 4 seconds.</p>	

Other Instructions:

Nurse _____

Responsible Adult _____