

What to Expect after Knee Arthroplasty

Andrea Saterbak, MD

- 1. You will spend 1 night in the hospital/care suite
- 2. You will be up and walking the day of surgery
 - a. A cane or walker will be provided by the facility for support
- **3.** Physical therapy (PT) should be done 2 times a week for at least 5 weeks and then down to 1 time a week until all goals are met
 - **a.** Regaining motion (flexion and extension), decreasing swelling, and pain management are the goals of therapy
- 4. The first 6 weeks after surgery are the most challenging!
 - **a.** There will be a lot of swelling above and below the knee (usually increases 3–5 days post-op)
 - i. Elevate leg above your heart
 - ii. Ice at least 5-6 times a day (20 minutes on 20 minutes off)
 - iii. Compression use stockings, ace wrap, or compression sleeve
 - iv. Ankle pumps (while leg is elevated, repeatedly point and flex foot)
 - Pushes swelling out of leg and back into the lymph system
 - v. Swelling will come and go throughout recovery
 - **b.** There will be pain (usually increases 3–4 days post-op)
 - i. Read through the Narcotics/Pain Medication handout
 - ii. This is part of the healing process
 - c. Range of motion will be limited due to pain and swelling
 - i. You have to push through pain to get motion back
- 5. Common post-surgical symptoms
 - a. Bruising above and below knee (entire leg may bruise)
 - i. Thigh bruising is due to the tourniquet used during surgery
 - **b. Post-surgical drainage on the bandage** this is normal and expected. Unless it is seeping out of the bandage, there is no need for concern.
 - c. Warmth to the touch this is normal
 - i. Symptoms of possible infections
 - Fever/chills
 - Drainage
 - Increase in pain different then surgical pain
 - Red streaks or increase in redness (usually bright red)
 - Unexplained loss of range of motion or the inability to walk
 - **d.** Swelling usually increases 3–5 days following surgery
 - e. Numbness

f. Stiffness

- i. Work through this in physical therapy and do at home stretches and exercises
- g. Trouble sleeping at night This is a common complaint!
 - i. Medications to try: Melatonin, Tylenol/Aleve
 - ii. Prednisone may be prescribed

6. Anticoagulant medications for blood clot prevention

- a. Aspirin 81mg 2/day for 6 weeks
- **b.** Other medications, as directed (if you are at risk for a blood clot)

7. Prescription pain medications

- a. There will be limited refills of the narcotic medications you may need to pay outof-pocket for refills due to insurance restrictions
- **b.** Read through *Narcotics/Prescription Medication* handout
- c. <u>Narcotic medications are used to help manage post-surgical pain, not to make it go away completely</u>
- **d.** Make sure to take stool softeners while using the narcotic medications

8. Showering:

- **a.** 3 days after surgery
- b. Let water run over large bandage
- c. Do not scrub
- **d.** Do not remove bandage
- e. Pat dry

9. Wound management:

- a. Keep large bandage on
- **b.** This will be removed at the first post-op appointment

10. Driving depends on the following:

- a. Which knee the surgery was on
- **b.** Must be off narcotic medications during the day
- **c.** Must be able to slam on the brakes
- **d.** If you feel you are ready, practice in a parking lot before you go on the roads

11. Precautions:

- a. Avoid pivoting, twisting, or kneeling on the operative leg
- **b.** Do not apply any heat to the knee or aggressively massage if for the first 2-3 weeks
- c. Avoid barefoot walking Wear supportive shoes/sandals to avoid stress on the knee
- **d.** Avoid submerging until 6-8 weeks after surgery
- e. Travel is not advised until about 8 weeks after surgery

*<u>Dental Visits</u>: No dental appointments for 3 months following a knee replacement. Following your knee replacement, you will need to be on an antibiotic prior to going to the dentist for 2 years. Call our office to get the prescription.

If you have any questions or concerns, please contact Dr. Saterbak's team at 651-351-2618 or SaterbakAcareteam@tcomn.com