

DISTAL CLAVICLE EXCISION (MUMFORD PROCEDURE) PT PROTOCOL

	BRACE INSTRUCTIONS AND RESTRICTIONS	RANGE OF MOTION GOALS	PT EXERCISES/NOTES
PHASE 1 First 4 Weeks (0-4) 1 Session/week or once every other week Begin after 2 weeks	 Sling x 2 weeks full time except hygiene/exercises, then part time during day x 2 weeks May remove sling at night after 2 weeks as tolerated Avoid horizontal adduction 	- PROM as tolerated in all positions	 First 2 weeks, begin Codman's After 2 weeks, progress posterior capsule mobilizations, closed chain scapular stabilizers
PHASE 2 Second 4 Weeks (4-8) 1-2 Sessions/week	 Out of sling Avoid horizontal adduction 	- Advance to full AAROM/AROM	 Wands/pulleys Wall climbs Initiate scapular stabilizing exercises, core strengthening exercises Begin resistive exercises for scapular stabilizers, biceps, triceps, and rotator cuff Begin rotator cuff strengthening after 6 weeks
PHASE 3 Third 4 Weeks (8-12) Sessions once/week or once every other week		- Full AROM in all planes	 Advance strength training as tolerated Begin eccentric resisted motion and closed chain activities Return to sport and full activity at 12 weeks

This protocol is intended for the purposes of guiding post-surgical or post-injury rehabilitation exercises and goals. These guidelines may be interpreted by a certified physical or occupation therapist and tailored to meet patient specific goals and expectations. Other specific modifications may be made by Dr. Butterfield depending on adjuvant surgical procedures, surgical findings, or rehabilitation progress.

Please fax initial physical therapy assessment and progress notes to Dr. Butterfield at 763-717-4139. You may direct questions regarding this protocol or patient progress to our office at 763-786-9543.