

ARTHROSCOPIC SLAP (SUPERIOR LABRUM ANTERIOR TO POSTERIOR) REPAIR PT PROTOCOL

	BRACE INSTRUCTIONS AND RESTRICTIONS	RANGE OF MOTION GOALS	PT EXERCISES/NOTES
PHASE 1 First 6 Weeks (0-6) 1-2 Sessions/week Begin after 2 weeks	<ul style="list-style-type: none"> - Shoulder immobilizer x 6 weeks full time except hygiene and PT exercises; may remove at night after 4 weeks 	<ul style="list-style-type: none"> - Begin PROM after 2 weeks (Goal of 135 deg FF, 120 deg Abduction, 45 deg ER by 6 weeks) - May begin AAROM at 4 weeks 	<ul style="list-style-type: none"> - Elbow/wrist ROM exercises, grip - Codman's, posterior capsule mobilizations, closed chain scapula exercises after 2 weeks
PHASE 2 Second 6 Weeks (6-12) 1-2 Sessions/week	<ul style="list-style-type: none"> - Out of brace - Avoid resisted rotator cuff exercises of the shoulder until 8 weeks - Avoid biceps resistive exercises until 8 weeks 	<ul style="list-style-type: none"> - Continue with PROM to reach full ROM and appropriate joint mobilization - Advance to AROM at 6 weeks 	<ul style="list-style-type: none"> - Wands/pulleys/wall climbs -Begin deltoid/rotator cuff isometrics at 6 weeks - Begin resistive exercises for scapular stabilizers, biceps, triceps, and rotator cuff at 8 weeks
PHASE 3 Third 6 Weeks (12-18) Sessions once/week or once every other week		<ul style="list-style-type: none"> - Full AROM in all planes 	<ul style="list-style-type: none"> - Progress PREs (light weight, high repetition) - Emphasize external rotation and latissimus eccentrics, dynamic stabilization - Neuromuscular and proprioceptive training - Aerobic conditioning and preparation for return to work/sport and ADLs - Cycling and running as tolerated

<p>PHASE 4 18 Weeks + Sessions as needed Transition to home exercise program</p>		<ul style="list-style-type: none"> - Sport and work specific exercises 	<ul style="list-style-type: none"> - Aggressive scapular stabilization and eccentric strengthening - Begin plyometric exercises - Begin throwing program if indicated - Improve neuromuscular control and shoulder proprioception - -Advance to home exercise program 3 times/week and daily shoulder stretching - -Return to sport and weight training after 4 months (contact sports after 5 months) once clear by MD
---	--	---	---

This protocol is intended for the purposes of guiding post-surgical or post-injury rehabilitation exercises and goals. These guidelines may be interpreted by a certified physical or occupation therapist and tailored to meet patient specific goals and expectations. Other specific modifications may be made by Dr. Butterfield depending on adjuvant surgical procedures, surgical findings, or rehabilitation progress.

Please fax initial physical therapy assessment and progress notes to Dr. Butterfield at 763-717-4139. You may direct questions regarding this protocol or patient progress to our office at 763-786-9543.