

**TJ Ridley, MD**

Phone: 763-717-4120 | Website: [TJRidleyMD.com](http://TJRidleyMD.com) | Instagram: @dr.\_ridley

**Clavicle Fracture Non-Operative Rehabilitation Protocol****PHASE 1: WEEK 0-4 (EARLY FRACTURE HEALING)**

- Maintain sling use (except for TID elbow wrist finger ROM, hygiene)
  - Elbow should be supported
- Rhomboid and periscapular isometrics, posture exercises
- No lifting > weight of coffee cup
- No active shoulder ROM
- No pushing or weight-bearing through arm

**PHASE 2: WEEK 4-8**

- Wean out of sling for therapy and in the house
  - Continue sling in public for 12 weeks or until clinical fracture healing
- Continue elbow, wrist and finger motion, periscapular isometrics
- Shoulder 4 quadrant PROM progressing towards AAROM
  - Pulleys, table slides, supine wand exercises in all planes
- No lifting > weight of coffee cup

**PHASE 3: WEEK 8-12 (FOLLOWING CLINICAL FRACTURE HEALING)**

- Continue sling in public for 12 weeks or until clinical fracture healing
- Advance to AROM
  - Continue 4 quadrant stretches, periscapular isometrics, posture
- Phase I and II cuff strengthening
- Periscapular and complete extremity strengthening
- No Lifting > 10lbs

**PHASE 4: WEEK 12-16**

- Discontinue sling altogether
- Independent home exercise program
- Return to sport
- Work hardening