

## TJ Ridley, MD

Phone: 763-717-4124 | Website: [TJRidleyMD.com](http://TJRidleyMD.com) | Instagram: @dr.\_ridley

### Proximal Humerus Fracture ORIF Rehabilitation Protocol

Phase / Goals	Immobilization	ROM	Therapeutic Exercises
<b>Phase 1 (0-2 weeks)</b> Goals: <ul style="list-style-type: none"> <li>Control pain and swelling</li> <li>Protect repair</li> </ul>	<ul style="list-style-type: none"> <li>Sling at all times x 3 weeks</li> </ul>	<ul style="list-style-type: none"> <li>PROM               <ul style="list-style-type: none"> <li>FF:0-90 deg</li> <li>ER:0-35 deg</li> <li>Abd:0-65 deg</li> </ul> </li> <li>Modalities PRN</li> <li>Soft tissue mobilization</li> </ul>	<ul style="list-style-type: none"> <li>Hand, wrist, elbow and cervical ROM</li> <li>Grip and wrist strengthening</li> <li>Pendulums (Codman)</li> <li>Scapular stabilization</li> </ul>
<b>Phase 2 (3-6 weeks)</b> Goals: <ul style="list-style-type: none"> <li>Frequent gentle exercises to prevent adhesions</li> <li>Increase ROM while protecting the fracture site</li> </ul>	<ul style="list-style-type: none"> <li>Wean from sling after week 3</li> </ul>	<ul style="list-style-type: none"> <li>PROM               <ul style="list-style-type: none"> <li>Full</li> </ul> </li> <li>AAROM               <ul style="list-style-type: none"> <li>Full</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>Isometrics FF and ER</li> <li>Pulley for flexion to tolerance</li> <li>Slide boards</li> <li>Scapular stabilization</li> </ul>
<b>Phase 3 (7-12 weeks)</b> Goals: <ul style="list-style-type: none"> <li>Full active ROM</li> <li>Muscular endurance</li> </ul>	<ul style="list-style-type: none"> <li>None</li> </ul>	<ul style="list-style-type: none"> <li>AROM               <ul style="list-style-type: none"> <li>Full</li> </ul> </li> <li>Begin gentle patient self-stretch</li> </ul>	<ul style="list-style-type: none"> <li>Multi-angle isometrics</li> <li>Advance to Theraband for IR, ER, FF and Abd</li> <li>Biceps and triceps strengthening</li> <li>Closed chain stability</li> </ul>

<b>Phase 4 (13-16 weeks)</b> Goals: <ul style="list-style-type: none"><li>• Maintain Full A/PROM</li><li>• Muscular strength</li></ul>	<ul style="list-style-type: none"><li>• None</li></ul>	<ul style="list-style-type: none"><li>• Full A/PROM</li><li>• Increase stretches</li><li>• Modalities PRN<ul style="list-style-type: none"><li>○ Deep tissue massage</li><li>○ Cupping</li><li>○ Graston</li></ul></li></ul>	<ul style="list-style-type: none"><li>• Increase Theraband in all directions</li><li>• Begin return to functional or sport exercises</li><li>• Final Xrays to demonstrate complete healing prior to return to sport</li></ul>
---	--	--	---