

TJ Ridley, MD

Phone: 763-717-4124 | Website: TJRidleyMD.com | Instagram: @dr._ridley

Total Shoulder Arthroplasty Rehabilitation Protocol

Phase / Goals	Immobilization	ROM	Therapeutic Exercises
Phase 1 (week 0-2) Goals: Pain and edema control PROM full FF and ABD, no ER > 30 x 2 weeks	Sling use full time x 4 weeks	PROM ER: 0-30 x 2 wks IR to belt FF: full ABD: full Modalities: IFC & ice x 20 mins	 Pendulums Cervical, elbow, wrist and hand ROM Strength: Wrist and grip only
Phase 2 (week 3-4) Goals: Passive ROM to limits outlined	Sling use full time x 4 weeks	PROM ER: 0-60 x 2 wks IR to belt FF: full ABD: full AAROM: within limits above	 Scapular retraction-depression Pendulums Begin isometrics wk 4 for ER No resisted elbow flexion x 6 wks Avoid reaching behind back x 6 wks
Phase 3 (week 5-6) Goals: • Full PROM by 6 week mark	D/C sling	Progress to full PROMBegin AROM	 Scapular retraction-depression Pendulums Isometrics Open chain proprioception Begin low load prolonged stretches No resisted elbow flexion x 6 wks

Phase 4 (week 7-12) Goals: • Full range of motion • 4/5 strength (rotator cuff and scapula stabilization)	• None	• Full A/PROM	Strength: Begin rotator cuff endurance Begin light therabands Posture control Aerobic: May start elliptical, treadmill at incline and progress to walk run x 30 mins Start UBE w/o resistance
Phase 5 (week 13-18) Goals: Full range of motion Full strength	• None	Continue full A/PROM	Aerobic: Continue progression to running on treadmill Rowing machine, Versiclimber Strength: Begin push-up progression Body blade at multiple angles Unrestricted therabands in all planes Continue rotator cuff strengthening Can return to full sport @ 6 month mark

CRITERIA FOR RETURN TO FULL ACTIVITY:

• Functional pain-free active range of motion, maximized strength, proper scapulothoracic mechanics

PRECAUTIONS:

• No repetitive lifting heavier than 50 lbs, no contact sports, only light-weight lifting