



**Kyle Flikkema, DO**

Phone: 763-786-9543 | Website: TCOmn.com/Kyle-Flikkema

**Knee Arthroscopic Meniscal Root and Radial Tear Repair Protocol**

Phase / Goals	Immobilization	ROM	Therapeutic Exercises
<b>Phase 1 (0-2 weeks)</b> Goals: <ul style="list-style-type: none"> <li>No quad lag</li> <li>Adequate hamstring control</li> </ul>	<ul style="list-style-type: none"> <li>Hinged brace 0-90</li> <li>Toe touch weight bearing x 6 weeks</li> </ul>	<ul style="list-style-type: none"> <li>0-90 deg A/PROM</li> <li>Patellar mobilization</li> <li>Modalities PRN               <ul style="list-style-type: none"> <li>Begin KneeHab</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>Wall slides, heel slides to 45 deg flex</li> <li>Quad activation, SLR, quad series</li> <li>Extension mobilization</li> <li>Sit &amp; reach for hamstring stretch w/ towel</li> <li>Ankle pumps</li> </ul>
<b>Phase 2 (3-6 weeks)</b> Goals: <ul style="list-style-type: none"> <li>Moderate proprioception</li> <li>Minimize swelling</li> </ul>	<ul style="list-style-type: none"> <li>Hinged brace 0-90</li> <li>TTWB x 6 weeks</li> </ul>	<ul style="list-style-type: none"> <li>0-90 deg A/PROM</li> <li>Patellar Mobilization</li> <li>Modalities PRN</li> </ul>	<ul style="list-style-type: none"> <li>Wall slides, heel slides</li> <li>Quad activation, SLR, quad series</li> <li>Extension mobilization</li> <li>Sit &amp; reach for hamstring stretch w/ towel</li> <li>Ankle pumps</li> <li>BFR – may initiate post op day 15</li> </ul>
<b>Phase 3 (7-10 weeks)</b> Goals: <ul style="list-style-type: none"> <li>Full ROM, advance to FWB with normalized gait</li> </ul>	<ul style="list-style-type: none"> <li>D/C hinged brace</li> <li>Advance 25% weekly weight bearing and progress to full WB with normalized gait</li> </ul>	<ul style="list-style-type: none"> <li>Advance to full A/PROM as tolerated</li> </ul>	<ul style="list-style-type: none"> <li>Can begin hamstring sets</li> <li>Toe &amp; heel raises</li> <li>Begin mini squats and progress as tolerated</li> <li>Begin Elliptical with minimal resistance</li> <li>Advance stationary bike with minimal resistance</li> <li>Aquajogging</li> </ul>
<b>Phase 4 (11-16 weeks)</b> Goals: <ul style="list-style-type: none"> <li>Return to golf, hiking, outdoor biking at 20 weeks</li> <li>Skiing, pivoting sports at 24 weeks</li> </ul>	<ul style="list-style-type: none"> <li>No brace</li> <li>FWB</li> </ul>	<ul style="list-style-type: none"> <li>Maintain full A/PROM</li> <li>No deep squats or sitting cross-legged x 4 months</li> </ul>	<ul style="list-style-type: none"> <li>Begin walk to run program</li> <li>Can begin rowing</li> <li>Leg press, single leg dead lift, balance squats</li> <li>Begin light plyometric training</li> <li>Sport specific drills at 16 weeks</li> <li>Clearance from Dr. Ridley prior to return to sport</li> </ul>

**RETURN-TO-SPORT CRITERIA:**

*Full return to all sports and games*

- At least 6 months from surgery
- No functional complaints, full range of motion & no effusion
- Confidence when running, cutting, jumping at full speed
- >90% isometric quadriceps strength
- >90% contralateral values on hop tests
- >90% Quad Index LSI with Biodex or HHD
- >90% Quad peak torque/body weight on Biodex (if applicable)